arXiv:1509.04529v1 [math.FA] 15 Sep 2015

LACUNARY STATISTICAL CONVERGENCE OF ORDER β IN DIFFERENCE SEQUENCES OF FUZZY NUMBERS

Hıfsı ALTINOK   Damla YAĞDIRAN
Department of Mathematics, Fırat University, 23119, Elazığ, TURKEY.
E-mail: hifsialtinok@gmail.com   dyagdiran@hotmail.com

Abstract. In this paper, we define the spaces $N_{\theta}^{\beta}\left(p,F,\Delta^{m}\right)$, $S_{\theta}^{\beta}\left(F,\Delta^{m}\right)$, $w_{p}^{\beta}\left(F,\Delta^{m}\right)$ for sequences of fuzzy numbers using generalized difference operator Δ^{m} and a lacunary sequence θ and give some relations between them, where $\beta\in(0,1]$ and $p>0$. Furthermore, in the last section of paper, some inclusion theorems are presented related to the spaces $S_{\theta}^{\beta}\left(F,\Delta^{m}\right)$ and $w_{p}^{\beta}\left(\theta,f,F,\Delta^{m}\right)$ according to modulus function f.

Key words and phrases: Fuzzy number, sequence of fuzzy numbers, statistical convergence, lacunary sequence, Cesàro summability, modulus function
Mathematics Subject Classification: 40A05; 40A25; 40A30; 40C05; 03E72.

1 Introduction

The idea of statistical convergence was given by Zygmund [26] in the first edition of his monograph published in Warsaw in 1935. The concept of statistical convergence was introduced by Steinhaus [22] and Fast [11] and later reintroduced by Schoenberg [21] independently. Over the years and under different names statistical convergence has been discussed in the theory of Fourier analysis, ergodic theory, number theory, measure theory, trigonometric series, turnpike theory and Banach spaces. Later on it was further investigated from the sequence space point of view and linked with summability theory by Connor [7], Et [8], Fridy [13], Mursaleen [17] and many others. In recent years, generalizations of statistical convergence have appeared in the study of strong integral summability and the structure of ideals of bounded continuous functions on locally compact spaces. Statistical convergence and its generalizations are also connected with subsets of the Stone-Čech compactification of the natural numbers. Moreover, statistical convergence is closely related to the concept of convergence in probability.

Now, we recall some basic concepts which we shall use throughout the paper.

By a lacunary sequence we mean an increasing integer sequence $\theta = (k_r)$ such that $k_0 = 0$ and $h_r = (k_r - k_{r-1}) \to \infty$ as $r \to \infty$. Throught this paper the intervals determined by θ will be denoted by $I_r = (k_{r-1}, k_r]$ and the ratio $\frac{k_r}{k_{r-1}}$ will be abbreviated by q_r. Recently lacunary sequence spaces have been studied in ([12],[14],[4])

The order of statistical convergence of a sequence of numbers was given by Gadjiev and Orhan [15] and later on statistical convergence of order α and strong $p-$Cesàro summability of order α was studied by Çolak [5]. Lacunary statistical convergence of order α in real number sequences was defined by Şengül and Et [23].

The difference spaces $\ell_\infty(\Delta)$, $c(\Delta)$ and $c_0(\Delta)$, consisting of all real valued sequences $x = (x_k)$ such that $\Delta x = \Delta^1 x = (x_k - x_{k+1})$ in the sequence spaces ℓ_∞, c and c_0, were defined by Kızmaz [16]. The idea of difference sequences was generalized by Et and Çolak [10], Altinok [2], Çolak *et al.* [6], Tripathy and Baruah [25] and many others.

A fuzzy number is a mapping $X : \mathbb{R} \longrightarrow [0,1]$ which satisfies the following properties:

i) X is normal, i.e., there exists an $t_0 \in \mathbb{R}$ such that $X(t_0) = 1$;

ii) X is fuzzy convex, i.e., for all $\lambda \in [0,1]$ and all $s, t \in \mathbb{R}$ and $0 \leq \lambda \leq 1, X(\lambda s + (1-\lambda)t) \geq \min[X(s), X(t)]$;

iii) X is upper semicontinuous on real numbers set;

iv) The set $\overline{\{t \in \mathbb{R} : X(t) > 0\}}$, denoted by $[X]^0$, is compact, where $\overline{\{t \in \mathbb{R} : X(t) > 0\}}$ is closure of $\{t \in \mathbb{R} : X(t) > 0\}$ with usual topology of $\mathbb{R}$.

$L(\mathbb{R})$ denotes the set of all fuzzy numbers on real numbers set $\mathbb{R}$ and is said to be a fuzzy number space.

For $\alpha \in (0,1]$, the α-level set $[X]^\alpha$ of fuzzy number X is defined by.

$$[X]^\alpha = \begin{cases} \{t \in \mathbb{R} : X(t) \geq \alpha\}, & \text{for } \alpha \in (0,1] \\ \overline{\{t \in \mathbb{R} : X(t) > \alpha\}}, & \text{for } \alpha = 0 \end{cases}$$

The aritmetic operations for α-level sets are defined as follows:

For $X, Y \in L(\mathbb{R})$, now and what follows we will denote $\alpha-$level sets by $[X]^\alpha = [u_1^\alpha, v_1^\alpha]$, $[Y]^\alpha = [u_2^\alpha, v_2^\alpha]$, $\alpha \in [0,1]$. Then we have

$$\begin{aligned} [X+Y]^\alpha &= [u_1^\alpha + u_2^\alpha, v_1^\alpha + v_2^\alpha] \\ [X-Y]^\alpha &= [u_1^\alpha - v_2^\alpha, v_1^\alpha - u_2^\alpha] \\ [X.Y]^\alpha &= \left[\min_{i,j\in[1,2]} u_i^\alpha v_j^\alpha, \max_{i,j\in[1,2]} u_i^\alpha v_j^\alpha\right]. \end{aligned}$$

A sequence $X = (X_k)$ of fuzzy numbers is a function X from the set $\mathbb{N}$ of all natural numbers into $L(\mathbb{R})$ [18].

Let $\beta \in (0,1]$ and $X = (X_k)$ be a sequence of fuzzy numbers. Then the sequence $X = (X_k)$ of fuzzy numbers is said to be statistically convergent of order β, to fuzzy number X_0 if for every $\varepsilon > 0$,

$$\lim_{n\to\infty} \frac{1}{n^{\beta}} \left|\{k \leq n : d\left(X_k, X_0\right) \geq \varepsilon\}\right| = 0,$$

where the vertical bars indicate the number of elements in the enclosed set. In this case we write $S^{\beta}\left(F\right) - \lim X_k = X_0$. We denote the set of all statistically convergent sequences of order β by $S^{\beta}\left(F\right)$ [3].

Let $w\left(F\right)$ be the set of all sequences of fuzzy numbers. The operator $\Delta^m : w\left(F\right) \to w\left(F\right)$ is defined by

$$\begin{aligned} \left(\Delta^0 X\right)_k &= X_k, \left(\Delta^1 X\right)_k = \Delta^1 X_k = X_k - X_{k+1}, \\ \left(\Delta^m X\right)_k &= \Delta^1 \left(\Delta^{m-1} X\right)_k, (m \geq 2), \text{ for all } k \in \mathbb{N}. \end{aligned}$$

2 Main Results

In this section, we define the spaces $S_{\theta}^{\beta}\left(F, \Delta^m\right)$ and $N_{\theta}^{\beta}\left(p, F, \Delta^m\right)$ and examine some inclusion relations between them and space $w_p^{\beta}\left(F, \Delta^m\right)$.

Definition 2.1 Let $\theta = (k_r)$ be a lacunary sequence, $X = (X_k)$ be a sequence of fuzzy numbers and $\beta \in (0,1]$ be given. The sequence $X = (X_k) \in w\left(F\right)$ is said to be $S_{\theta}^{\beta}\left(F, \Delta^m\right)$ −statistically convergent (or lacunary statistically convergent sequence of order β) if there is a fuzzy number X_0 such that

$$\lim_{r\to\infty} \frac{1}{h_r^{\beta}} \left|\{k \in I_r : d\left(\Delta^m X_k, X_0\right) \geq \varepsilon\}\right| = 0,$$

where $I_r = (k_{r-1}, k_r]$ and h_r^{β} denote the βth power $(h_r)^{\beta}$ of h_r, that is $h^{\beta} = (h_r^{\beta}) = (h_1^{\beta}, h_2^{\beta}, ..., h_r^{\beta}, ...)$. In this case we write $S_{\theta}^{\beta}\left(F, \Delta^m\right) - \lim X_k = X_0$. The set of all $S_{\theta}^{\beta}\left(F, \Delta^m\right)$ −statistically convergent sequences will be denoted by $S_{\theta}^{\beta}\left(F, \Delta^m\right)$. For $\theta = (2^r)$, we shall write $S^{\beta}\left(F, \Delta^m\right)$ instead of $S_{\theta}^{\beta}\left(F, \Delta^m\right)$ and we shall write $S\left(F, \Delta^m\right)$ instead of $S_{\theta}^{\beta}\left(F, \Delta^m\right)$ in the special case $\beta = 1$ and $\theta = (2^r)$.

Lemma 2.2. Let $\theta = (k_r)$ be a lacunary sequence and $K \subset \mathbb{N}$. Then $S_{\theta}^{\beta}\left(K\right) \leq S_{\theta}^{\alpha}\left(K\right)$ for $0 < \alpha \leq \beta \leq 1$.

Proof. Let $0 < \alpha \leq \beta \leq 1$. We can write

$$\frac{1}{h_r^{\beta}}\left|\left\{k_{r-1}<k\leq k_r : k\in K\right\}\right| \leq \frac{1}{h_r^{\alpha}}\left|\left\{k_{r-1}<k\leq k_r : k\in K\right\}\right|$$

since $h_r^{\alpha}\leq h_r^{\beta}$. Hence $S_{\theta}^{\beta}(K)\leq S_{\theta}^{\alpha}(K)$.

Let $\beta\in(0,1]$. If a sequence of fuzzy numbers is lacunary statistically convergent of order β, then it is lacunary statistically convergent. Really, we obtain

$$\frac{1}{h_r^{\alpha}}\left|\left\{k\in I_r : d\left(\Delta^m X_k, X_0\right)\geq\varepsilon\right\}\right| \geq \frac{1}{h_r}\left|\left\{k\in I_r : d\left(\Delta^m X_k, X_0\right)\geq\varepsilon\right\}\right|$$

since $h_r^{\alpha}\leq h_r$ for $\beta\in(0,1]$. Taking limit as $r\to\infty$ we get $X_k\to X_0\left(S_{\theta}^{\alpha},\Delta^m\right)$ and so $X_k\to X_0\left(S_{\theta},\Delta^m\right)$.

Note that the lacunary statistical convergence of order β is well defined for $\beta\in(0,1]$, but not well defined for $\beta>1$. To show this, consider the sequence $X=(X_k)$ of fuzzy numbers following:

$$X_k(t)=\begin{cases}\left.\begin{array}{cc} t, & \text{for } 0\leq t\leq 1\\ -t+2, & \text{for } 1\leq t\leq 2\\ 0, & \text{otherwise}\end{array}\right\} := X', & \begin{array}{c}\text{if } k=2r\\ (r=1,2,3,...)\end{array}\\ \left.\begin{array}{cc} t-3, & \text{for } 3\leq t\leq 4\\ -t+5, & \text{for } 4\leq t\leq 5\\ 0, & \text{otherwise}\end{array}\right\} := X'', & \text{if } k\neq 2r\end{cases}$$

We can find the $\alpha-$level sets of sequences (X_k) and $(\Delta^m X_k)$ after some arithmetic operations as follows

$$[X_k]^{\alpha}=\begin{cases}[\alpha,2-\alpha], & \text{if } k=2r\\ [3+\alpha,5-\alpha], & \text{if } k\neq 2r\end{cases}.$$

and

$$[\Delta^m X_k]^{\alpha}=\begin{cases}\left[2^m\alpha-2^{m-1}.5,-2^m\alpha-2^{m-1}\right] := X', & \text{if } k=2r\\ \left[2^m\alpha+2^{m-1},-2^m\alpha+2^{m-1}.5\right] := X'', & \text{if } k\neq 2r\end{cases},(m=1,2,...)$$

then we can write

$$\lim_{r\to\infty}\frac{1}{h_r^{\beta}}\left|\left\{k\in I_r : d\left(\Delta^m X_k, X'\right)\geq\varepsilon\right\}\right| \leq \lim_{r\to\infty}\frac{k_r-k_{r-1}}{2h_r^{\beta}}=\lim_{r\to\infty}\frac{h_r}{2h_r^{\beta}}=0$$

and

$$\lim_{r\to\infty}\frac{1}{h_r^{\beta}}\left|\left\{k\in I_r : d\left(\Delta^m X_k, X''\right)\geq\varepsilon\right\}\right| \leq \lim_{r\to\infty}\frac{k_r-k_{r-1}}{2h_r^{\beta}}=\lim_{r\to\infty}\frac{h_r}{2h_r^{\beta}}=0$$

for $\beta > 1$. Hence sequence (X_k) is Δ^m-lacunary statistically convergent of order β, both to X' and X'', i.e., $S_\theta^\beta(F,\Delta^m) - \lim X_k = X'$ and $S_\theta^\beta(F,\Delta^m) - \lim X_k = X''$. But this is impossible.

Theorem 2.3 Let $0 < \beta \leq 1$ and $X = (X_k)$, $Y = (Y_k)$ be sequences of fuzzy numbers, then

(*i*) If $S_\theta^\beta(F,\Delta^m) - \lim X_k = X_0$ and $c \in \mathbb{C}$, then $S_\theta^\beta(F,\Delta^m) - \lim(cx_k) = cx_0$,

(*ii*) If $S_\theta^\beta(F,\Delta^m) - \lim X_k = X_0$ and $S_\theta^\beta(F,\Delta^m) - \lim Y_k = Y_0$, then $S_\theta^\beta(F,\Delta^m) - \lim(X_k + Y_k) = X_0 + Y_0$.

Proof. (*i*) It is clear for the case $c = 0$. Suppose that $c \neq 0$, then the proof of (*i*) follows from

$$\frac{1}{h_r^\beta}\left|\{k \in I_r : d(c\Delta^m X_k, cX_0) \geq \varepsilon\}\right| = \frac{1}{h_r^\beta}\left|\left\{k \in I_r : d(\Delta^m X_k, X_0) \geq \frac{\varepsilon}{|c|}\right\}\right|$$

and that of (*ii*) follows from

$$\begin{aligned}\frac{1}{h_r^\beta}\left|\{k \in I_r : d(\Delta^m(X_k+Y_k),(X_0+Y_0)) \geq \varepsilon\}\right| \;\leq\; & \frac{1}{h_r^\beta}\left|\left\{k \in I_r : d(\Delta^m X_k, X_0) \geq \frac{\varepsilon}{2}\right\}\right| \\ & + \frac{1}{h_r^\beta}\left|\left\{k \in I_r : d(\Delta^m Y_k, Y_0) \geq \frac{\varepsilon}{2}\right\}\right|.\end{aligned}$$

Definition 2.4 Let $X = (X_k)$ be a sequence of fuzzy numbers, $\theta = (k_r)$ be a lacunary sequence, $\beta \in (0,1]$ be any real number and let p be a positive real number. A sequence X of fuzzy numbers is said to be strongly $N_\theta^\beta(p,F,\Delta^m)-$summable (or strongly $N_\theta(p,F,\Delta^m)-$summable of order β) if there is a fuzzy number X_0 such that

$$\lim_{r\to\infty}\frac{1}{h_r^\beta}\sum_{k\in I_r} d(\Delta^m X_k, X_0)^p = 0.$$

In this case we write $N_\theta^\beta(p,F,\Delta^m) - \lim X_k = X_0$. The strong $N_\theta^\beta(p,F,\Delta^m)-$summability reduces to the strong $N_\theta(p,F,\Delta^m)-$summability for $\beta = 1$. The set of all strongly $N_\theta^\beta(p,F,\Delta^m)-$summable sequences will be denoted by $N_\theta^\beta(p,F,\Delta^m)$.

If we take $\theta = (2^r)$ in space $N_\theta^\beta(p,F,\Delta^m)$, then we obtain strongly $p-$Cesaro summable sequences set of order β following

$$w_p^\beta(F,\Delta^m) = \left\{X = (X_k) : \exists X_0, \lim_n \frac{1}{n^\beta}\sum_{k=1}^n [d(\Delta^m X_k, X_0)]^p = 0\right\}$$

This set will be denoted by $w_{p,0}^\beta$ in case of $X_0 = \bar{0}$ [2].

Theorem 2.5 If $0 < \beta < \gamma \leq 1$ then $S_\theta^\beta(F,\Delta^m) \subset S_\theta^\gamma(F,\Delta^m)$ and the inclusion is strict.

Proof. Proof can be obtained from the inequality

$$\frac{1}{h_r^{\gamma}}\left|\left\{k\in I_r: d\left(\Delta^m X_k, X_0\right)\geq\varepsilon\right\}\right|\leq\frac{1}{h_r^{\beta}}\left|\left\{k\in I_r: d\left(\Delta^m X_k, X_0\right)\geq\varepsilon\right\}\right|.$$

Taking $\theta=(2^r)$ we show the strictness of the inclusion $S_\theta^\beta(F,\Delta^m)\subset S_\theta^\gamma(F,\Delta^m)$ for a special case. For this, consider the sequence $X=(X_k)$ of fuzzy numbers defined by

$$X_k(t)=\begin{cases}\left.\begin{array}{cc}2t-(2k-1), & \text{for } k-\frac{1}{2}\leq t\leq k\\ -2t+(2k+1), & \text{for } k\leq t\leq k+\frac{1}{2}\\ 0, & \text{otherwise}\end{array}\right\}, & \text{if } k=n^3\\ \left.\begin{array}{cc}2t-1, & \text{for } \frac{1}{2}\leq t\leq 1\\ -2t+3, & \text{for } 1\leq t\leq\frac{3}{2}\\ 0, & \text{otherwise}\end{array}\right\}, & \text{if } k\neq n^3\end{cases}$$

After calculating $\alpha-$level sets of sequences (X_k) and $(\Delta^m X_k)$, we obtain $S_\theta^\gamma(F,\Delta^m)-\lim X_k=\left[2^{m-1}(\alpha-1),2^{m-1}(1-\alpha)\right]$, i.e. $X\in S_\theta^\gamma(F,\Delta^m)$ for $\frac{1}{3}<\gamma\leq 1$, but $X\notin S_\theta^\beta(F,\Delta^m)$ for $0<\beta\leq\frac{1}{3}$. This implies that the inclusion $S_\theta^\beta(F,\Delta^m)\subset S_\theta^\gamma(F,\Delta^m)$ is strict for $\beta,\gamma\in(0,1]$ such that $\beta\in\left(0,\frac{1}{3}\right]$ and $\gamma\in\left(\frac{1}{3},1\right]$.

Corollary 2.6 If a sequence of fuzzy numbers is $S_\theta^\beta(F,\Delta^m)-$statistically convergent to fuzzy number X_0, then it is $S_\theta(F,\Delta^m)-$statistically convergent to fuzzy number X_0.

Theorem 2.7 Let β and γ be fixed real numbers such that $0<\beta\leq\gamma\leq 1$, $X=(X_k)$ be a sequence of fuzzy numbers and $\theta=(k_r)$ be a lacunary sequence. For $0<p<\infty$, $N_\theta^\beta(p,F,\Delta^m)\subset S_\theta^\gamma(F,\Delta^m)$ and the inclusion is strict for some β's and γ's.

Proof. For any sequence $X=(X_k)$ of fuzzy numbers and $\varepsilon>0$, we have

$$\begin{aligned}\sum_{k\in I_r}\left[d\left(\Delta^m X_k,X_0\right)\right]^p &= \sum_{\substack{k\in I_r\\ d(\Delta^m X_k,X_0)\geq\varepsilon}}\left[d\left(\Delta^m X_k,X_0\right)\right]^p+\sum_{\substack{k\in I_r\\ d(\Delta^m X_k,X_0)<\varepsilon}}\left[d\left(\Delta^m X_k,X_0\right)\right]^p\\ &\geq \sum_{\substack{k\in I_r\\ d(\Delta^m X_k,X_0)\geq\varepsilon}}\left[d\left(\Delta^m X_k,X_0\right)\right]^p\\ &\geq \left|\left\{k\in I_r: d\left(\Delta^m X_k,X_0\right)\geq\varepsilon\right\}\right|\varepsilon^p\end{aligned}$$

and so that

$$\begin{aligned}\frac{1}{h_r^\beta}\sum_{k\in I_r}\left[d\left(\Delta^m X_k,X_0\right)\right]^p &\geq \frac{1}{h_r^\beta}\left|\left\{k\in I_r: d\left(\Delta^m X_k,X_0\right)\geq\varepsilon\right\}\right|\varepsilon^p\\ &\geq \frac{1}{h_r^\gamma}\left|\left\{k\in I_r: d\left(\Delta^m X_k,X_0\right)\geq\varepsilon\right\}\right|\varepsilon^p.\end{aligned}$$

It follows that if $X=(X_k)$ is strongly $N_\theta^\beta(p,F,\Delta^m)-$summable to fuzzy number X_0, then it is $S_\theta^\gamma(F,\Delta^m)-$statistically convergent to X_0.

Taking $\beta=\gamma$, $\theta=(2^r)$ and $p=1$ we show the strictness of the inclusion $N_\theta^\beta(p,F,\Delta^m)\subset S_\theta^\gamma(F,\Delta^m)$ for a special case. For this we can select a sequence of fuzzy numbers as follows:

$$X_k(t)=\begin{cases}\left.\begin{array}{cc}\frac{t}{k}+1, & \text{for } -k\le t\le 0\\ -\frac{t}{k}+1, & \text{for } 0\le t\le k\\ 0, & \text{otherwise}\end{array}\right\}, & \begin{array}{c}\text{if } k=n^2\\ (n=1,2,...)\end{array}\\ \left.\begin{array}{cc}\frac{t}{2}-1, & \text{for } 2\le t\le 4\\ -\frac{t}{2}+3, & \text{for } 4\le t\le 6\\ 0, & \text{otherwise}\end{array}\right\}:=X_0 & \text{if } k\ne n^2\end{cases}$$

After calculating $\alpha-$level sets of sequences (X_k) and $(\Delta^m X_k)$, we see that (X_k) is Δ^m-statistically convergent of order γ to X_0 for $\frac{1}{2}<\beta\le 1$, where $[X_0]^\alpha=\left[2^{m+1}(\alpha-1),2^m(1-\alpha)\right]$, but it is not strongly $N_\theta^\beta(p,F,\Delta^m)$-summable to X_0. Therefore $N_\theta^\beta(p,F,\Delta^m)\subset S_\theta^\beta(F,\Delta^m)$ for $\frac{1}{2}<\beta\le 1$.

Corollary 2.8 Let $X=(X_k)$ be a sequence of fuzzy numbers, $\beta\in(0,1]$ and p is a positive real number. Then

i) If $X\in N_\theta^\beta(p,F,\Delta^m)$, then $X\in S_\theta^\beta(F,\Delta^m)$ and the limits are same.

ii) If $X\in N_\theta^\beta(p,F,\Delta^m)$, then $X\in S_\theta(F,\Delta^m)$ and the limits are same.

Theorem 2.9 Let $X=(X_k)$ be a sequence of fuzzy numbers, $\beta\in(0,1]$ and $\theta=(k_r)$ be a lacunary sequence. If $\liminf_r q_r>1$, then $S^\beta(F,\Delta^m)\subset S_\theta^\beta(F,\Delta^m)$.

Proof. Suppose that $\liminf_r q_r>1$, then there exists a number $\delta>0$ such that $q_r\ge 1+\delta$ for sufficiently large r and so

$$\frac{h_r}{k_r}\ge\frac{\delta}{1+\delta}\Longrightarrow\left(\frac{h_r}{k_r}\right)^\beta\ge\left(\frac{\delta}{1+\delta}\right)^\beta\Longrightarrow\frac{1}{k_r^\beta}\ge\frac{\delta^\beta}{(1+\delta)^\beta}\frac{1}{h_r^\beta}.$$

If $X_k\to X_0\left(S^\beta(F,\Delta^m)\right)$, then we prove the sufficiency from following inequalities for every $\varepsilon>0$ and for sufficiently large r

$$\begin{aligned}\frac{1}{k_r^\beta}\left|\{k\le k_r:d(\Delta^m X_k,X_0)\ge\varepsilon\}\right| &\ge \frac{1}{k_r^\beta}\left|\{k\in I_r:d(\Delta^m X_k,X_0)\ge\varepsilon\}\right|\\ &\ge \frac{\delta^\beta}{(1+\delta)^\beta}\frac{1}{h_r^\beta}\left|\{k\in I_r:d(\Delta^m X_k,X_0)\ge\varepsilon\}\right|;\end{aligned}$$

Theorem 2.10 If $\lim_{r\to\infty}\inf\frac{h_r^\beta}{k_r}>0$, then $S(F,\Delta^m)\subset S_\theta^\beta(F,\Delta^m)$.

Proof. For a given $\varepsilon > 0$, we can write

$$\{k \leq k_r : d\left(\Delta^m X_k, X_0\right) \geq \varepsilon\} \supset \{k \in I_r : d\left(\Delta^m X_k, X_0\right) \geq \varepsilon\}.$$

From here we obtain

$$\begin{aligned} \frac{1}{k_r}\left|\{k \leq k_r : d\left(\Delta^m X_k, X_0\right) \geq \varepsilon\}\right| &\geq \frac{1}{k_r}\left|\{k \in I_r : d\left(\Delta^m X_k, X_0\right) \geq \varepsilon\}\right| \\ &= \frac{h_r^\beta}{k_r}\frac{1}{h_r^\beta}\left|\{k \in I_r : d\left(\Delta^m X_k, X_0\right) \geq \varepsilon\}\right|. \end{aligned}$$

Taking limit according to r and using $\lim_{r\to\infty} \inf \frac{h_r^\beta}{k_r} > 0$, we get

$$X_k \to X_0\left(S, F, \Delta^m\right) \Longrightarrow X_k \to X_0\left(S_\theta^\beta, F, \Delta^m\right).$$

Theorem 2.11 Let $X = (X_k)$ be a sequence of fuzzy numbers, $0 < \beta \leq 1$,and $\theta = (k_r)$ be a lacunary sequence. If $\limsup_r q_r < \infty$, then $S_\theta^\beta\left(F, \Delta^m\right) \subset S\left(F, \Delta^m\right)$.

Proof. Omitted.

Theorem 2.12 Let $0 < \beta < \gamma \leq 1$, $X = (X_k)$ be a sequence of fuzzy numbers and $p > 0$, then there exists inclusion $N_\theta^\beta\left(p, F, \Delta^m\right) \subset N_\theta^\gamma\left(p, F, \Delta^m\right)$ and this is strict.

Proof. Let $X = (X_k) \in N_\theta^\beta\left(p, F, \Delta^m\right)$. Then given β and γ such that $0 < \beta < \gamma \leq 1$ and a $p > 0$, we can write inequality

$$\frac{1}{h_r^\gamma}\sum_{k\in I_r}\left[d\left(\Delta^m X_k, X_0\right)\right]^p \leq \frac{1}{h_r^\beta}\sum_{k\in I_r}\left[d\left(\Delta^m X_k, X_0\right)\right]^p$$

and this gives that $N_\theta^\beta\left(p, F, \Delta^m\right) \subset N_\theta^\gamma\left(p, F, \Delta^m\right)$.

To show that the inclusion is strict take lacunary sequence $\theta = (2^r)$ and $p = 1$. Consider the sequence $X = (X_k)$ of fuzzy numbers as follows

$$X_k(t) = \left\{\begin{array}{ll} \left.\begin{array}{cc} t-2, & \text{for } 2 \leq t \leq 3 \\ -t+4, & \text{for } 3 \leq t \leq 4 \\ 0, & \text{otherwise} \end{array}\right\}, & \begin{array}{c} \text{if } k = n^3 \\ (n = 1, 2, ...) \end{array} \\ \left.\begin{array}{cc} \frac{t}{3}-\frac{5}{3}, & \text{for } 5 \leq t \leq 8 \\ -\frac{t}{3}+\frac{11}{3}, & \text{for } 8 \leq t \leq 11 \\ 0, & \text{otherwise} \end{array}\right\} := X_0 & \text{if } k \neq n^3 \end{array}\right.$$

After calculating $\alpha-$level sets of sequences (X_k) and $(\Delta^m X_k)$, we obtain $X \in N_\theta^\gamma\left(p, F, \Delta^m\right)$ since (X_k) is strongly $N_\theta^\beta\left(p, F, \Delta^m\right)$-summable to X_0 for $\frac{1}{3} < \gamma \leq 1$, where $[X_0]^\alpha = \left[2^m\left(3\alpha - 3\right), 2^m\left(3 - 3\alpha\right)\right]$, but $X \notin N_\theta^\beta\left(p, F, \Delta^m\right)$ for $0 < \beta \leq \frac{1}{3}$, i.e. it is not strongly $N_{\theta}^{\beta}(p,F,\Delta^m)$-summable to fuzzy number X_0. So, this shows that inclusion is strict. We show the limits of sequence $\Delta^m X_k$ in Figure 1.

Corollary 2.13 Let $0<\beta\leq\gamma\leq 1$, $X=(X_k)$ be a sequence of fuzzy numbers and $p>0$. Then

(*i*) If $\beta=\gamma$, then $N_{\theta}^{\beta}(p,F,\Delta^m)=N_{\theta}^{\gamma}(p,F,\Delta^m)$,

(*ii*) For each $\beta\in(0,1]$ and $0<p<\infty$, $N_{\theta}^{\beta}(p,F,\Delta^m)\subseteq N_{\theta}(p,F,\Delta^m)$.

Theorem 2.14. Let $X=(X_k)$ be a sequence of fuzzy numbers and $\theta=(k_r)$ be a lacunary sequence. If $\liminf_r q_r>1$, then $w_p^{\beta}(F,\Delta^m)\subseteq N_{\theta}^{\beta}(p,F,\Delta^m)$ for $0<\beta\leq 1$ and $0<p<\infty$.

Proof. If $\liminf_r q_r>1$, then there exists a number $\delta>0$ such that $q_r\geq 1+\delta$ for all $r\geq 1$. For $0<\beta\leq 1$ and $X\in w_{p,0}^{\beta}$ we can write

$$\begin{aligned}
\tau_r^{\beta} &= \frac{1}{h_r^{\beta}}\sum_{i=1}^{k_r} d(\Delta^m X_i,\bar{0})^p-\frac{1}{h_r^{\beta}}\sum_{i=1}^{k_{r-1}} d(\Delta^m X_i,\bar{0})^p\\
&= \frac{k_r^{\beta}}{h_r^{\beta}}\left(\frac{1}{k_r^{\beta}}\sum_{i=1}^{k_r} d(\Delta^m X_i,\bar{0})^p\right)-\frac{k_{r-1}^{\beta}}{h_r^{\beta}}\left(\frac{1}{k_{r-1}^{\beta}}\sum_{i=1}^{k_{r-1}} d(\Delta^m X_i,\bar{0})^p\right)
\end{aligned}$$

and so

$$\frac{k_r^{\beta}}{h_r^{\beta}}\leq\frac{(1+\delta)^{\beta}}{\delta^{\beta}}\text{ and }\frac{k_{r-1}^{\beta}}{h_r^{\beta}}\leq\frac{1}{\delta^{\beta}}$$

for $h_r=k_r-k_{r-1}$. It can be seen that both of terms $\frac{1}{k_r^{\beta}}\sum_{i=1}^{k_r} d(\Delta^m X_i,\bar{0})^p$ and $\frac{1}{k_{r-1}^{\beta}}\sum_{i=1}^{k_{r-1}} d(\Delta^m X_i,\bar{0})^p$ converge to 0 and so τ_r^{β} does. Hence $X\in N_{\theta,0}^{\beta}(p,F,\Delta^m)$, that is $w_p^{\beta}(F,\Delta^m)\subseteq N_{\theta}^{\beta}(p,F,\Delta^m)$.

Theorem 2.15. If $\limsup_r \dfrac{k_r}{k_{r-1}^{\beta}} < \infty$, then $N_\theta\left(p, F, \Delta^m\right) \subseteq w_p^{\beta}\left(F, \Delta^m\right)$ for $0 < \beta \leq 1$.

Proof. If $\limsup_r \dfrac{k_r}{k_{r-1}^{\beta}} < \infty$, then there exists a number $M > 0$ such that $\dfrac{k_r}{k_{r-1}^{\beta}} < M$ for all $r \geq 1$. Let $X \in N_{\theta,0}\left(p, F, \Delta^m\right)$ and $\varepsilon > 0$. Then we can find numbers $R > 0$ and $K > 0$ such that $\sup_{i \geq R} \tau_i < \varepsilon$ and $\tau_i < K$ for $i = 1, 2, 3, ...$. Therefore we can write

$$\begin{aligned}
\frac{1}{t^{\beta}} \sum_{i=1}^{t} d\left(\Delta^m X_i, \bar{0}\right)^p &\leq \frac{1}{k_{r-1}^{\beta}} \sum_{i=1}^{k_{r-1}} d\left(\Delta^m X_i, \bar{0}\right)^p \\
&= \frac{1}{k_{r-1}^{\beta}} \left(\sum_{I_1} d\left(\Delta^m X_i, \bar{0}\right)^p + \sum_{I_2} d\left(\Delta^m X_i, \bar{0}\right)^p + ... + \sum_{I_r} d\left(\Delta^m X_i, \bar{0}\right)^p \right) \\
&= \frac{k_1}{k_{r-1}^{\beta}} \tau_1 + \frac{k_2 - k_1}{k_{r-1}^{\beta}} \tau_2 + ... + \frac{k_R - k_{R-1}}{k_{r-1}^{\beta}} \tau_R + \frac{k_{R+1} - k_R}{k_{r-1}^{\beta}} \tau_{R+1} + ... + \frac{kr - k_{r-1}}{k_{r-1}^{\beta}} \tau_r \\
&\leq \left(\sup_{i \geq 1} \tau_i \right) \frac{k_R}{k_{r-1}^{\beta}} + \left(\sup_{i \geq R} \tau_i \right) \frac{k_r - k_R}{k_{r-1}^{\beta}} < K \frac{k_R}{k_{r-1}^{\beta}} + \epsilon M
\end{aligned}$$

where $k_{r-1} < t \leq k_r$ and $r > R$. We get $\frac{1}{t^{\beta}} \sum_{i=1}^{t} d\left(\Delta^m X_i, \bar{0}\right)^p \to 0$ since $t \to \infty$ for $k_{r-1} \to \infty$, hence $X \in w_{p,0}^{\beta}\left(F, \Delta^m\right)$.

Theorem 2.16. Let $X = (X_k)$ be a sequence of fuzzy numbers, $\theta = (k_r)$ be a lacunary sequence and $\beta \in (0, 1]$. If $X \in w^{\beta}\left(F, \Delta^m\right) \cap N_\theta^{\beta}\left(p, F, \Delta^m\right)$ and $\limsup_r \dfrac{k_r}{k_{r-1}^{\beta}} < \infty$, then the limits of (X_k) are same.

Proof. Suppose that $w_p^{\beta}\left(F, \Delta^m\right) - \lim X_k = X'$, $N_{\theta,p}^{\beta}\left(p, F, \Delta^m\right) - \lim X_k = X''$ and $X' \neq X''$. We get $N_{\theta,0}\left(p, F, \Delta^m\right) \subseteq w_{p,0}^{\beta}\left(F, \Delta^m\right)$ since $\limsup_r \dfrac{k_r}{k_{r-1}^{\beta}} < \infty$ from Theorem 2.15. Because of $d\left(\Delta^m X_i, X''\right) \in N_{\theta,0}\left(p, F, \Delta^m\right)$, so $d\left(\Delta^m X_i, X''\right) \in w_{p,0}^{\beta}\left(F, \Delta^m\right)$. Therefore for $p = 1$

$$\frac{1}{t^{\beta}} \sum_{i=1}^{t} d\left(\Delta^m X_i, X''\right) \to 0$$

In the following inequality, both of terms at left converges to 0, but this is a contradiction and so $X' = X''$.

$$\frac{1}{t^{\beta}}\sum_{i=1}^{t} d\left(\Delta^m X_i, X''\right)+\frac{1}{t^{\beta}}\sum_{i=1}^{t} d\left(\Delta^m X_i, X'\right)\geq \frac{1}{t^{\beta}} d\left(X', X''\right)>0$$

3. Results Related to Modulus Function

We give relation between sets $S_{\theta}^{\beta}\left(F,\Delta^m\right)$ and $w_p^{\beta}\left[\theta, f, F, \Delta^m\right]$ according to modulus function f.

We first quote the definition of a modulus function:

The concept of modulus function was formally introduced by Nakano [19]. A mapping $f:[0,\infty)\rightarrow[0,\infty)$ is said to be a modulus if

i) $f(x)=0$ iff $x=0$,

ii) $f(x+y)\leq f(x)+f(y)$ for $x, y\geq 0$,

iii) f is increasing,

iv) f is right-continuous at 0.

The continuity of f everywhere on $[0,\infty)$ follows from above definition. A modulus function can be bounded or unbounded. For example $f(x)=x^p, (0<p\leq 1)$ is bounded and $f(x)=\frac{x}{1+x}$ is bounded. The concept of modulus function for sequences of fuzzy numbers was first investigated by Sarma [20]. Later on it was explotied by Talo and Başar [24].

Definition 3.1. Let f be a modulus function, $X=(X_k)$ be a sequence of fuzzy numbers and $p=(p_k)$ be a sequence of positive real numbers and $\beta\in(0,1]$. We define space $w_p^{\beta}\left(\theta, f, F, \Delta^m\right)$ as follows

$$w_p^{\beta}\left(\theta, f, F, \Delta^m\right)=\left\{x=(x_k): \lim_{r\rightarrow\infty}\frac{1}{h_r^{\beta}}\sum_{k\in I_r}\left[d\left(\Delta^m X_k, X_0\right)\right]^{p_k}=0,\ \text{for some } X_0\right\}$$

We shall write $N_{\theta}^{\beta}\left(p, F, \Delta^m\right)$ instead of $w_p^{\beta}\left(\theta, f, F, \Delta^m\right)$ in the special cases $p_k=1$ and $f(x)=x$ for all $k\in\mathbb{N}$.

We consider that $p=(p_k)$ is bounded and $0<h=\inf_k p_k\leq p_k\leq \sup_k p_k=H<\infty$ in the following theorems.

Theorem 3.2. Let $\beta,\gamma\in(0,1]$, $\beta\leq\gamma$, f be a modulus function, $X=(X_k)$ be a sequence of fuzzy numbers and $\theta=(k_r)$ be a lacunary sequence. Then, $w_p^{\beta}\left(\theta, f, F, \Delta^m\right)\subset S_{\theta}^{\gamma}\left(F,\Delta^m\right).$

Proof. Take $x \in w_p^\beta(\theta, f, F, \Delta^m)$, $\varepsilon > 0$ and let $\sum\limits_1$ and $\sum\limits_2$ be summations on $k \in I_r$, $d(\Delta^m X_k, X_0) \geq \varepsilon$ and $k \in I_r$, $d(\Delta^m X_k, X_0) < \varepsilon$, respectively. Since $h_r^\beta < h_r^\gamma$ for each r, we get

$$\begin{aligned}
\frac{1}{h_r^\beta} \sum_{k \in I_r} [f(d(\Delta^m X_k, X_0))]^{p_k} &= \frac{1}{h_r^\beta} \left[\sum_1 [f(d(\Delta^m X_k, X_0))]^{p_k} + \sum_2 [f(d(\Delta^m X_k, X_0))]^{p_k} \right] \\
&\geq \frac{1}{h_r^\gamma} \left[\sum_1 [f(d(\Delta^m X_k, X_0))]^{p_k} + \sum_2 [f(d(\Delta^m X_k, X_0))]^{p_k} \right] \\
&\geq \frac{1}{h_r^\gamma} \sum_1 [f(\varepsilon)]^{p_k} \\
&\geq \frac{1}{h_r^\gamma} \sum_1 \min\left([f(\varepsilon)]^h, [f(\varepsilon)]^H \right) \\
&\geq \frac{1}{h_r^\gamma} \left| \{k \in I_r : d(\Delta^m X_k, X_0) \geq \varepsilon\} \right| \min\left([f(\varepsilon)]^h, [f(\varepsilon)]^H \right)
\end{aligned}$$

Left side of above inequality converges to 0 as $r \to \infty$ because of $X \in w_p^\beta(\theta, f, F, \Delta^m)$, so right one converges to zero and hence $X \in S_\theta^\gamma(F, \Delta^m)$.

Theorem 3.3. Let f be a bounded modulus function, $X = (X_k)$ be a sequence of fuzzy numbers and $\theta = (k_r)$ be a lacunary sequence. If $\lim\limits_{r \to \infty} \dfrac{h_r}{h_r^\beta} = 1$, then $S_\theta^\beta(F, \Delta^m) \subset w_p^\beta(\theta, f, F, \Delta^m)$.

Proof. Suppose that f is bounded and $X \in S_\theta^\beta(F, \Delta^m)$. Given $\varepsilon > 0$. There is an integer K such that $(x) \leq K$ since f is bounded. Now we can write

$$\begin{aligned}
\frac{1}{h_r^\beta} \sum_{k \in I_r} [f(d(\Delta^m X_k, X_0))]^{p_k} &= \frac{1}{h_r^\beta} \sum_1 [f(d(\Delta^m X_k, X_0))]^{p_k} + \frac{1}{h_r^\beta} \sum_2 [f(d(\Delta^m X_k, X_0))]^{p_k} \\
&\leq \frac{1}{h_r^\beta} \sum_1 \max\left(K^h, K^H \right) + \frac{1}{h_r^\beta} \sum_2 [f(\varepsilon)]^{p_k} \\
&\leq \max\left(K^h, K^H \right) \frac{1}{h_r^\beta} \left| \{k \in I_r : f(d(\Delta^m X_k, X_0)) \geq \varepsilon\} \right| \\
&\quad + \frac{h_r}{h_r^\beta} \max\left(f(\varepsilon)^h, \ f(\varepsilon)^H \right).
\end{aligned}$$

Hence we get $X \in w_p^\beta(\theta, f, F, \Delta^m)$ for $r \in \mathbb{N}$ from above inequalities.

Theorem 3.4. If $\lim p_k > 0$ and sequence $X = (X_k)$ is strongly $w_p^\beta(\theta, f, F, \Delta^m)$ $-$summable to X_0, then $w_p^\beta(\theta, f, F, \Delta^m) - \lim X_k = X_0$ is unique.

Proof. Let $\lim p_k = s > 0$, $w_p^\beta(\theta, f, F, \Delta^m) - \lim X_k = X'$ and $w_p^\beta(\theta, f, F, \Delta^m) -$ $\lim X_k = X''$. Then we obtain

$$\lim_{r\to\infty} \frac{1}{h_r^\beta} \sum_{k\in I_r} \left[f\left(d\left(\Delta^m X_k, X'\right)\right)\right]^{p_k} = 0$$

and

$$\lim_{r\to\infty} \frac{1}{h_r^\beta} \sum_{k\in I_r} \left[f\left(d\left(\Delta^m X_k, X''\right)\right)\right]^{p_k} = 0$$

On the other hand, since f is increasing and $f(x+y) \leq f(x) + f(y)$ we can write

$$\begin{aligned} \frac{1}{h_r^\beta} \sum_{k\in I_r} \left[f\left(d\left(X', X''\right)\right)\right]^{p_k} &\leq \frac{D}{h_r^\beta} \sum_{k\in I_r} \left(\left[f\left(d\left(\Delta^m X_k, X'\right)\right)\right]^{p_k} + \left[f\left(d\left(\Delta^m X_k, X''\right)\right)\right]^{p_k}\right) \\ &\leq \frac{D}{h_r^\beta} \sum_{k\in I_r} \left[f\left(d\left(\Delta^m X_k, X'\right)\right)\right]^{p_k} + \frac{D}{h_r^\beta} \sum_{k\in I_r} \left[f\left(d\left(\Delta^m X_k, X''\right)\right)\right]^{p_k} \to 0, \end{aligned}$$

where $p_k = H$ and $D = \left(1, 2^{H-1}\right)$. Left right of above inequalities tends to zero as $r \to \infty$ and because of $\lim p_k = s$ we obtain

$$\lim_{r\to\infty} \left[f\left(d\left(X', X''\right)\right)\right]^{p_k} = \left[f\left(d\left(X', X''\right)\right)\right]^s$$

Hence $X' = X''$ and the limit is unique.

References

[1] Altin, Y. ; Et, M. and Tripathy, B. C. The sequence space $|\bar{N}, p_k|(M, r, q, s)$ on seminormed spaces, Appl. Math. Comput. 154(2) (2004), 423–430.

[2] Altınok, H. Statistical convergence of order β for generalized difference sequences of fuzzy numbers, Journal of Intelligent & Fuzzy Systems, 26, 847–856, 2014.

[3] Altınok, H. Altın, Y. and Işık, M. Statistical Convergence and Strong $p-$Cesàro Summability of Order β in Sequences of Fuzzy Numbers, Iranian J. of Fuzzy Systems, 9 (2), 65-75, 2012

[4] Altınok, H. Et, M. and Çolak, R. Some remarks on generalized sequence space of bounded variation of sequences of fuzzy numbers, Iranian J. of Fuzzy Systems, 11(5), pp. 39-46, 2014.

[5] Çolak, R. Statistical convergence of order α, Modern Methods in Analysis and Its Applications, New Delhi, India: Anamaya Pub, 2010: 121–129.

[6] Çolak, R., Altınok, H. and Et, M. Generalized difference sequences of fuzzy numbers, Chaos, Solitons & Fractals 40, 1106–1117, 2009.

[7] Connor, J. S. The statistical and strong $p-$Cesaro convergence of sequences, Analysis 8 (1988), 47-63.

[8] Et, M. Strongly almost summable difference sequences of order m defined by a modulus, Studia Sci. Math. Hungar. 40(4) (2003), 463–476.

[9] Et. M. Spaces of Cesàro difference sequences of order r defined by a modulus function in a locally convex space, Taiwanese J. Math. 10(4) (2006), 865–879.

[10] Et, M. and Çolak, R. On some generalized difference sequence spaces, Soochow J. Math. 21 (4), 377-386, 1995

[11] Fast, H. Sur la convergence statistique, Colloq. Math. 2 (1951), 241-244.

[12] Freedman, A. R. Sember J. J. and Raphael, M. Some Cesaro-type summability spaces, Proc. Lond. Math. Soc. 37(3) (1978), 508-520.

[13] Fridy, J. On statistical convergence, Analysis 5 (1985), 301-313.

[14] Fridy J.A. and Orhan, C. Lacunary statistical convergence, Pacific J. Math. 160 (1993), 43-51.

[15] Gadjiev, A. D. and Orhan, C. Some approximation theorems via statistical convergence, Rocky Mountain J. Math. 32(1) (2002), 129-138.

[16] Kızmaz, H. On certain sequence spaces, Canad. Math. Bull. 24 (2), 169-176, 1981.

[17] Mursaleen, M. $\lambda-$ statistical convergence, Math. Slovaca, 50(1) (2000), 111 -115.

[18] Matloka, M. Sequences of fuzzy numbers, BUSEFAL 28 (1986), 28-37.

[19] Nakano, H. Concave modulars, J. Math. Soc. Japan, 5, (1953), 29–49.

[20] Sarma, B. On a class of sequences of fuzzy numbers defined by modulus function, International Journal of Science & Technology, 2(1), (2007), 25-28.

[21] Schoenberg, I. J. The integrability of certain functions and related summability methods, Amer. Math. Monthly 66 (1959), 361-375.

[22] Steinhaus, H. Sur la convergence ordinaire et la convergence asymptotique, Colloquium Mathematicum 2 (1951),73-74.

[23] Şengül, H. and Et, M. On lacunary statistical convergence of order α, Acta Mathematica Scientia, 34B (2), 473-482, 2014.

[24] Talo, Ö. and Başar, F. Certain spaces of sequences of fuzzy numbers defined by a modulus function, Demonstratio Math. 43 (2010), no. 1, 139–149.

[25] Tripathy, B.C. and Baruah, A. Lacunary statistically convergent and lacunary strongly convergent generalized difference sequences of fuzzy real numbers, Kyungpook Math. Jour. 50, 565-574, 2010.

[26] Zygmund, A. Trigonometric Series, Cambridge University Press, Cambridge, UK, 1979.